

## APPENDIX B

Calculations of Site-Specific Background Concentrations of Metals in Soil



#### **APPENDIX B**

# SITE-SPECIFIC BACKGROUND CONCENTRATIONS OF METALS IN SOIL

Former Pechiney Cast Plate, Inc. Facility 3200 Fruitland Avenue Vernon, California

#### 1.0 INTRODUCTION

Naturally-occurring inorganic constituents (i.e., metals) detected in soil at a site may be eliminated as chemicals of potential concern (COPCs) if detected concentrations are consistent with local or site-specific background conditions (also referred to as "ambient conditions," or conditions unaffected by past site-related activities). The derivation of local or site-specific background concentrations may also be used to evaluate if remedial action or risk management measures specific to metals in soil is warranted. Because of the high density of industrial land use surrounding the former Pechiney Cast Plate, Inc. Facility (the Site), collecting soil samples to establish local background concentrations would not be appropriate or applicable to the Site. Instead, site-specific background concentrations of metals in soil were derived for the Site from on-site data as described herein. The analysis presented has been prepared in accordance with several California Environmental Protection Agency (Cal-EPA) Department of Toxic Substances Control (DTSC) guidance documents including:

- Selecting Inorganic Constituents as Chemicals of Potential Concern at Risk Assessments at Hazardous Waste Sites and Permitted Facilities, February 1997 (DTSC, 1997);<sup>1</sup> and
- Arsenic Strategies, Determination of Arsenic Remediation, Development of Arsenic Cleanup Goals, January 16, 2009 (DTSC, 2009).<sup>2</sup>

Information regarding the lithology of the Site, along with a description of the previous on-site investigations for metals in soil, is presented in Sections 2.0 through 3.0 of the Feasibility Study (FS). The metals data collected from these previous investigations and evaluated for use in the derivation of site-specific background concentrations are presented in Appendix A

<sup>&</sup>lt;sup>1</sup> Department of Toxic Substances Control (DTSC), 1997, Selecting Inorganic Constituents as Chemicals of Potential Concern at Risk Assessment at Hazardous Waste Sites and Permitted Facilities, February.

<sup>&</sup>lt;sup>2</sup> Department of Toxic Substances Control (DTSC), 2009, Arsenic Strategies, Determination of Arsenic Remediation, Development Arsenic Cleanup Goals, January 16.



of the FS. A description of the approach and methods used to derive the site-specific background concentrations is presented in the following sections.

#### 2.0 DATA EVALUATION

As presented in Appendix A of the FS, 486 soil samples were collected from the ground surface to a depth of 50 feet below ground surface (bgs) at the Site between 1991 and 2007 and analyzed for metals using U.S. EPA Method 6010B or U.S. EPA Method 7471A (for mercury). Of these 486 soil samples, 249 soil samples are no longer in place following excavations (marked as "Excavated" in Appendix A). Only the analytical results from the 237 "left-in-place" soil samples were considered for the evaluation of background conditions. Summary statistics for the analytical results of these samples are presented in Table B-1.

Metals with low frequency of detection were excluded from the evaluation of site-specific background. Specifically, antimony, beryllium, cadmium, hexavalent chromium, molybdenum, selenium, silver, and thallium were excluded, with detection frequencies between 0 and 6 percent. The detection frequencies for the remaining 10 metals detected at the Site (arsenic, barium, total chromium, cobalt, copper, lead, mercury, nickel, vanadium, and zinc) were considered sufficient to warrant statistical and graphical evaluation.

Detection limits were not available for two non-detect results of lead for samples UST-2B-1 and UST-2B-2 collected in 1995. The non-detect results were simply listed as ND in the provided data tables associated with the removal of the underground storage tanks (laboratory reports for these samples were not available). These non-detect results were therefore excluded from the subsequent statistical evaluations.

#### 3.0 SITE-SPECIFIC BACKGROUND CONCENTRATIONS

The statistical and graphical methods applied to evaluate and identify site-specific background concentrations for metals in soil included a goodness-of-fit and probability plots. First, each metal was evaluated using goodness-of-fit tests and probability plots to determine if single or multiple data populations exist within each dataset. Gaps or inflection points identified from the probability plots, for example, would be indicative of a shift from background concentrations to site-related concentrations (DTSC, 1997). For those metals for which inflection points could not be identified but for which the goodness-of-fit tests suggested multiple data populations were present, additional statistical evaluations were performed to identify outliers (impacted soil samples) and estimate site-specific background concentrations.



#### 3.1 DISTRIBUTION EVALUATION AND IDENTIFICATION OF INFLECTION POINTS

Consistent with DTSC guidance (DTSC, 1997), the distribution of each dataset was evaluated to determine if multiple distributions were present. Distributions were tested using the Lilliefor's goodness-of-fit test function in U.S. EPA's ProUCL product, Version 4.00.04 (ProUCL) (U.S. EPA, 2009)<sup>3</sup> and were also evaluated graphically with normal and lognormal probability plots generated using ProUCL. Dataset distributions that fail goodness-of-fit tests for normality and lognormality and/or the presence of inflection points in a probability plot generally indicate that a dataset is comprised of multiple populations (DTSC, 1997). In these cases, background conditions are defined as the range of concentrations associated with the population nearest the origin (i.e., the first population) (DTSC, 1997; 2009).

The results of the goodness-of-fit testing for each metal are presented in Attachment B-1. As shown, only one dataset was considered to fit a normal, lognormal, or gamma theoretical distribution; the distribution of the mercury data set was determined to fit a lognormal distribution, but only when excluding the non-detect results. Therefore, it is likely that the datasets of all 10 metals evaluated are comprised of both background and site-related data populations.

Normal and lognormal probability plots are presented for each metal in Attachment B-2. Detection limits ranged widely from samples collected and analyzed during the 1990s to more recent analytical results. For example, the detection limits for non-detect results of arsenic ranged from 0.05 milligrams per kilogram (mg/kg) to 60 mg/kg (for two samples, H-1 and H-2; Appendix A). The presence of multiple non-detect results within the range of detected concentrations created "noise" within some of the probability plots. To aid in the identification of distribution types and possible gaps or inflection points, normal and lognormal probability plots were also prepared by excluding these non-detect results.

Inflection points were identified for the following metals from the various probability plots (Attachment B-2):

- Arsenic Inflection point at 9.9 mg/kg, based on the normal probability plot with nondetect results removed;
- Copper Inflection point at 35 mg/kg, based on the normal probability plot of the complete data set;
- Mercury Inflection point at 0.18 mg/kg, based on the normal or lognormal probability plots with non-detect results removed;



- Nickel Inflection point at 17 mg/kg, based on the normal probability plots for the complete data set or with the non-detect results removed; and
- Zinc Inflection point at 92 mg/kg, based on the normal probability plot of the complete data set.

These inflection points were identified as the site-specific background concentrations for these metals, with one exception. As the inflection point for arsenic, 9.9 mg/kg, was in agreement with a 10 mg/kg background concentration previously established by the City of Vernon using comparable graphical methods (City of Vernon H&EC, letter dated April 28, 2008),<sup>4</sup> 10 mg/kg was used as the site-specific background concentration for arsenic at the Site.

Inflection points were not identified from the probability plots for barium, total chromium, cobalt, lead, or vanadium that would potentially distinguish background from site-related data populations. As a result, the datasets for these metals were subjected to further statistical evaluations to estimate site-specific background concentrations as described in Section 3.2.

#### 3.2 OUTLIER EVALUATION AND ESTIMATION OF UPPER LIMITS

Because inflection points could not be identified from the probability plots for barium, total chromium, cobalt, lead, and vanadium, upper limit concentrations were quantitatively estimated as representative of site-specific background concentrations. Each dataset was first evaluated for potential outliers. Outliers should be removed prior to estimating an upper limit concentration to ensure the upper limit estimate is not overly influenced by one or two observations, but "conforms to the pattern established by the majority of values in the dataset" (DTSC, 2009). Rosner's test was applied with ProUCL to qualitatively test for the presence of outliers (see Attachment B-3). Outliers were identified in the total chromium, lead, and vanadium background datasets and were excluded from their respective datasets prior to estimating the upper limit concentrations. The outliers identified by Rosner's test are depicted in the probability plots in Attachment B-2.

<sup>&</sup>lt;sup>3</sup> U.S. EPA, 2009, ProUCL Version 4.00.04, Technical Guide, Office of Research and Development, Washington DC, April.

<sup>&</sup>lt;sup>4</sup> City of Vernon Health & Environmental Control, 2008, letter to Ms. Linda Conlan re: Comments on Revised Geomatrix Feasibility Study/Remedial Action Plan (FS/RAP), Former Pechiney Cast Plate, Inc. Facility, 3200 Fruitland Avenue, Vernon, California, April 28.



With the suspected outliers removed, the upper limit concentrations for barium, total chromium, cobalt, lead, and vanadium were estimated using the following equation (DTSC, 2009):

$$UL = x bar + K * sd$$
 (1)

Where: UL = the upper limit of the dataset

x\_bar = the mean of detected concentrations in the dataset

K = statistical tolerance factor for estimating an upper confidence

limit on a given percentile of the population

\* = multiply

sd = the standard deviation of the mean

The value of the statistical tolerance factor was determined from a table in Gilbert (1987)<sup>5</sup> based on sample size, upper confidence limit, and quantile of the data population. As defined by DTSC (DTSC, 2009), the 95 percent upper confidence limit on the 99<sup>th</sup> quantile of each dataset was estimated for each metal, using a statistical tolerance factor based on data population parameters. The calculation of the upper limit concentrations of background is presented in Table B-2 for barium, total chromium, cobalt, lead, and vanadium.

#### 3.3 SUMMARY OF SITE-SPECIFIC BACKGROUND CONCENTRATIONS

A summary of the site-specific background concentrations for metals at the Site, identified as inflection points from normality plots or quantified as upper limit concentrations from background populations, is presented in Table B-3.

-

<sup>&</sup>lt;sup>5</sup> Gilbert, R.O., 1987, *Statistical Methods for Environmental Pollution Monitoring*. Van Nostrand Reinhold Co., New York.



## TABLE B-1

#### SUMMARY STATISTICS OF ANALYTICAL RESULTS FOR METALS

Former Pechiney Cast Plate, Inc. Facility Vernon, California

Results are shown in milligrams per kilogram (mg/kg)

						. 1000.10 0.	C 3110WIT III III	iiigiaiiie pe	(	9,9/								
						,	Chromium,											
Sample Date	Antimony	Arsenic	Barium	Beryllium	Cadmium	Hexavalent	Total	Cobalt	Copper	Lead	Mercury	Molybdenum	Nickel	Selenium	Silver	Thallium	Vanadium	Zinc
Count	217	216	217	217	217	23	219	217	219	221	217	217	217	217	217	217	217	217
Number of Detects	0	147	217	0	13	1	219	203	219	186	61	2	208	2	1	4	217	217
Number of Non-Detects	217	69	0	217	204	22	0	14	0	35	156	215	9	215	216	213	0	0
Mean Detected Concentration	NA	5.0	92.3	NA	1.25	0.35	12.9	8.0	20.3	9.4	0.12	6	9.6	1.41	5	1.24	32.5	58.0
Median Concentration	NA	1.9	93.9	NA	NA	NA	13	8	17	2.8	NA	NA	9.3	NA	NA	NA	34.0	51.0
Standard Deviation of Detections	NA	13.0	33.1	NA	0.66	NA	4.8	2.4	23.6	18.7	0.14	1.41	3.5	1.12	NA	0.61	9.6	52.1
Coefficient of Variation of Detections	NA	2.6	0.4	NA	0.53	NA	0.37	0.30	1.2	2.0	1.21	0.24	0.36	0.79	NA	0.49	0.30	0.9
Minimum Detection	0	0.63	23	0	0.54	0.35	2.7	2.2	3.3	0.62	0.023	5	2.4	0.62	5	0.74	7.6	13.3
Maximum Detection	0	120	190	0	2.8	0.35	32.1	16	257	157	0.98	7	27	2.2	5	2.1	70	607
Minimum Reporting Limit	5	0.05	0	0.5	0.5	0.04	0	3	0	3	0.02	4	3	0.5	1	0.5	0	0
Maximum Reporting Limit	12	60	0	1	1	0.04	0	10	0	20	0.1	8	4	1	2	100	0	0
Detection Frequency	0%	68%	100%	0%	6%	4%	100%	94%	100%	84%	28%	1%	96%	1%	0.5%	2%	100%	100%

Abbreviations:

NA = not applicable



**TABLE B-2** 

#### **CALCULATION OF UPPER LIMIT CONCENTRATIONS**

Former Pechiney Cast Plate, Inc. Facility Vernon, California

Results are shown in milligrams per kilogram (mg/kg)

	Barium	Chromium, Total	Cobalt	Lead	Vanadium
First Population					
Number of Detections	217	218	203	185	216
Number of Outliers	0	1 (32.1)	0	1 (157)	1 (70)
Mean Detection	92.3	12.9	7.95	8.59	32.3
Standard Deviation of Detections	33.1	4.63	2.35	15.3	9.25
Κ	2.62	2.62	2.62	2.62	2.62
UL <sub>95</sub> (X <sub>99</sub> )	179	25	14.1	48.5	56.5

#### Abbreviations:

K = statistical tolerance factor (from Gilbert, 1987).  $UL_{95}(X_{99}) = 95$  percent upper limit for the 99th quantile.

#### Equations:

$$UL_{95}(X_{99}) = x_bar + K * sd$$



#### **TABLE B-3**

#### SUMMARY OF SITE-SPECIFIC BACKGROUND CONCENTRATIONS FOR METALS

Former Pechiney Cast Plate, Inc. Facility Vernon, California

Results are shown in milligrams per kilogram (mg/kg)

Metal	Site-Specific Background Concentration	Basis
Arsenic	10	Inflection Point <sup>1</sup>
Barium	179	Upper Limit Concentration
Chromium, Total	25	Upper Limit Concentration
Cobalt	14.1	Upper Limit Concentration
Copper	35	Inflection Point
Lead	48.5	Upper Limit Concentration
Mercury	0.18	Inflection Point
Nickel	17	Inflection Point
Vanadium	56.5	Upper Limit Concentration
Zinc	92	Inflection Point

#### Notes:

<sup>1.</sup> As the inflection point for arsenic, 9.9 mg/kg, was in agreement with a 10 mg/kg background concentration previously established by the City of Vernon using comparable graphical methods (City of Vernon H&EC, letter dated April 28, 2008); 10 mg/kg was used as the site-specific background concentration for arsenic at the Site.

## ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS



	v emon,	California				
rsenic						
	Num Obs	Num Miss	Num Valid	Dotooto	NDs	% ND
Raw Statistics	216	0	216	Detects 147	69	0.319
Naw Statistics	210	O	210	147	03	0.515
	Number	Minimum	Maximum	Mean	Median	SD
Statistics (Non-Detects Only)	69	0.05	60	15.92	1	16.5
Statistics (Detects Only)	147	0.63	120	4.982	2.5	13.01
Statistics (All: NDs treated as DL value)	216	0.05	120	8.477	2.5	15.08
Statistics (All: NDs treated as DL/2 value)	216	0.025	120	5.934	2.5	11.77
Statistics (Normal ROS Estimated Data)	216	-21.89	120	1.761	2.15	12.74
Statistics (Gamma ROS Estimated Data)	216	1E-09	120	4.507	2.35	11.19
Statistics (Lognormal ROS Estimated Data)	216	0.185	120	3.878	2.103	10.89
	K Hat	K Star	Theta Hat	Log Mean	Log Stdv	Log C
Statistics (Detects Only)	0.947	0.937	5.263	0.992	0.789	0.79
Statistics (NDs = DL)	0.598	0.593	14.17	1.104	1.364	1.236
Statistics (NDs = DL/2)	0.676	0.67	8.777	0.882	1.364	1.546
Statistics (Gamma ROS Estimates)	0.173	0.174	26.03			
Statistics (Lognormal ROS Estimates)				0.671	0.944	1.40
No		* T+ D				
Noi	mai Distribu	tion Test Res	ults			
	No NDs	NDs = DL	NDs = DL/2	Normal ROS		
Correlation Coefficient R	0.491	0.728	0.646	0.728		
	Test value	Crit. (0.05)	Conclusion v	vith Alpha(0.0	5)	
Lilliefors (Detects Only)	0.414	0.0731	Data Not No	rmal		
Lilliefors (NDs = DL)	0.378	0.0603	Data Not No	rmal		
Lilliefors (NDs = DL/2)	0.324	0.0603	Data Not No	rmal		
Lilliefors (Normal ROS Estimates)	0.287	0.0603	Data Not No	rmal		
Gar	nma Distribu	ıtion Test Res	ults			
	No NDs	NDs = DL	NDs = DL/2	Gamma ROS	2	
Correlation Coefficient R	0.729	0.938	0.879	0.925	•	
Correlation Coemicient it	0.723	0.550	0.073	0.323		
	Test value	Crit. (0.05)	Conclusion v	vith Alpha(0.0	5)	
Anderson-Darling (Detects Only)	20.47	0.786	Conclusion	vitii Aipiia(0.0		
Kolmogorov-Smirnov (Detects Only)	0.308	0.0798	Data Not Ga	mma Distribu	ted	
Anderson-Darling (NDs = DL)	18.73	0.811	Data Not Ga			
Kolmogorov-Smirnov (NDs = DL)	0.277	0.0649	Data Not Ga	mma Distribu	ted	
Anderson-Darling (NDs = DL/2)	9.118	0.804		a Diombu		
Kolmogorov-Smirnov (NDs = DL/2)	0.208	0.0646	Data Not Ga	mma Distribu	ted	
Anderson-Darling (Gamma ROS Estimates)	40.75	0.968		a Diombu		
Kolmogorov-Smirnov (Gamma ROS Est.)	0.382	0.0698	Data Not Ga	mma Distribu	ted	
		3.2.200			-	
Logn	ormal Distrib	oution Test Re	esults			
	No NDs	NDs = DL	NDs = DL/2	Log ROS		
Correlation Coefficient R	0.895	0.952	0.97	0.963		
	Test value	Crit. (0.05)	Conclusion v	vith Alpha(0.0	15)	
Lilliefors (Detects Only)	0.187	0.0731	Data Not Log		*	
Lilliefors (NDs = DL)	0.174	0.0603	Data Not Log			
	•					
Lilliefors (NDs = DL/2)	0.125	0.0603	Data Not Log	normal		

#### ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS



	Num Obe	Num Miss	Num Valid	Detects	NDs	% NE
Raw Statistics 2:		20	217	217	0	0.00%
Naw Glatistics 25	57	20	217	217	O	0.0070
	Number	Minimum	Maximum	Mean	Median	SD
Statistics (Full: no NDs) 2	17	23	190	92.28	93.9	33.06
	K Hat	K Star	Theta Hat	Log Mean	Log Stdv	Log C
Statistics (Full: no NDs) 6.	.173	6.091	14.95	4.442	0.445	0.1
Norm	nal Distribut	tion Test Res	ults			
	No NDs	NDs = DL	NDs = DL/2	Normal ROS	3	
Correlation Coefficient R 0.	.992	0.992	0.992	0.992		
7	Test value	Crit. (0.05)		Conclusion w	rith Alpha(0.0	5)
Lilliefors (Full: no NDs) 0.	.0772	0.0601	Data Not No	ormal		
Gamr	ma Distribu	tion Test Res	ults			
	No NDs	NDs = DL	NDs = DL/2	Gamma RO	S	
Correlation Coefficient R 0.	.968	0.968	0.968	0.968		
1	Test value	Crit. (0.05)		Conclusion w	rith Alpha(0.0	5)
Anderson-Darling (Full: no NDs) 4.	.447	0.755				
Kolmogorov-Smirnov (Full: no NDs) 0.	.112	0.0618	Data Not Ga	amma Distribi	uted	
Lognor	mal Distrib	ution Test Re	sults			
	No NDs	NDs = DL	NDs = DL/2	Log ROS		
Correlation Coefficient R 0.	.947	0.947	0.947	0.947		
1	Test value	Crit. (0.05)		Conclusion w	rith Alpha(0.0	5)
Lilliefors (Full: no NDs) 0.	.14	0.0601	Data Not Lo	gnormal		

#### ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS



	Num Obs	Num Miss	Num Valid	Detects	NDs	% NI
Raw Statistics 2	237	18	219	219	0	0.00%
	Number	Minimum	Maximum	Mean	Median	SE
Statistics (Full: no NDs) 2	219	2.7	32.1	12.95	13	4.797
	K Hat	K Star	Theta Hat	Log Mean	Log Stdv	Log (
Statistics (Full: no NDs) 5	5.491	5.419	2.358	2.467	0.481	0.195
Nom	nal Distribut	tion Test Res	ults			
	No NDs	NDs = DL	NDs = DL/2	Normal ROS	3	
Correlation Coefficient R 0	).981	0.981	0.981	0.981		
	Test value	Crit. (0.05)		Conclusion w	rith Alpha(0.0	5)
Lilliefors (Full: no NDs) 0	).102	0.0599	Data Not No	ormal		
Gam	ma Distribu	tion Test Res	ults			
	No NDs	NDs = DL	NDs = DL/2	Gamma RO	S	
Correlation Coefficient R 0	).955	0.955	0.955	0.955		
	Test value	Crit. (0.05)		Conclusion w	rith Alpha(0.0	5)
Anderson-Darling (Full: no NDs) 7	7.381	0.756				
Kolmogorov-Smirnov (Full: no NDs) 0	).167	0.0616	Data Not Ga	amma Distribi	uted	
Logno	rmal Distrib	ution Test Re	sults			
	No NDs	NDs = DL	NDs = DL/2	Log ROS		
Correlation Coefficient R 0	).928	0.928	0.928	0.928		
	Test value	Crit. (0.05)		Conclusion w	rith Alpha(0.0	5)
Lilliefors (Full: no NDs) 0	).195	0.0599	Data Not Lo	gnormal		

## ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS



	Num Obs	Num Miss	Num Valid	Detects	NDs	9
Raw Statistics		20	217	203	14	6.4
	Number	Minimum	Maximum	Mean	Median	
Statistics (Non-Detects Only)	14	3	10	5.286	5	2.16
Statistics (Detects Only)	203	2.2	16	7.951	8.1	2.3
Statistics (All: NDs treated as DL value)	217	2.2	16	7.779	8	2.42
Statistics (All: NDs treated as DL/2 value)	217	1.5	16	7.609	8	2.63
Statistics (Normal ROS Estimated Data)	217	2.2	16	7.717	8	2.46
Statistics (Gamma ROS Estimated Data)	217	1.57	16	7.722	8	2.48
Statistics (Lognormal ROS Estimated Data)	217	2.2	16	7.727	8	2.4
	K Hat	K Star	Theta Hat	Log Mean	Log Stdv	L
Statistics (Detects Only)	8.802	8.684	0.903	2.015	0.373	0.1
Statistics (NDs = DL)	8.115	8.006	0.959	1.989	0.386	0.19
Statistics (NDs = DL/2)	6.014	5.934	1.265	1.944	0.462	0.2
Statistics (Gamma ROS Estimates)	7.371	7.272	1.048			
Statistics (Lognormal ROS Estimates)				1.981	0.388	0.1
No	rmal Distribu	tion Test Res	ults			
	No NDs	NDs = DL	NDs = DL/2	Normal ROS	3	
Correlation Coefficient R	0.981	0.985	0.976	0.984		
	Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)
Lilliefors (Detects Only)	0.1	0.0622	Data Not No	rmal		
Lilliefors (NDs = DL)	0.0945	0.0601	Data Not No	rmal		
Lilliefors (NDs = DL/2)	0.117	0.0601	Data Not No	rmal		
Lilliefors (Normal ROS Estimates)	0.101	0.0601	Data Not No	rmal		
Ga	mma Distribu	ition Test Res	ults			
	No NDs	NDs = DL	NDs = DL/2	Gamma RO	S	
Correlation Coefficient R	0.956	0.961	0.94	0.955		
	Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)
Anderson-Darling (Detects Only)	6.774	0.752				
Kolmogorov-Smirnov (Detects Only)	0.155	0.0633	Data Not Ga	ımma Distrib	uted	
Anderson-Darling (NDs = DL)	6.133	0.753				
Kolmogorov-Smirnov (NDs = DL)	0.148	0.0617	Data Not Ga	ımma Distrib	uted	
Anderson-Darling (NDs = DL/2)	9.8	0.755				
Kolmogorov-Smirnov (NDs = DL/2)	0.178	0.0618	Data Not Ga	ımma Distrib	uted	
Anderson-Darling (Gamma ROS Estimates)	7.216	0.754				
Kolmogorov-Smirnov (Gamma ROS Est.)	0.157	0.0617	Data Not Ga	ımma Distrib	uted	
Logr	normal Distrib	oution Test Re	esults			
	No NDs	NDs = DL	NDs = DL/2	Log ROS		
Correlation Coefficient R	0.92	0.935	0.912	0.939		
	Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)
Lilliefors (Detects Only)	0.18	0.0622	Data Not Lo	gnormal		
	0.172	0.0601	Data Not Lo	gnormal		
Lilliefors (NDs = DL)	0.172					
Lilliefors (NDs = DL) Lilliefors (NDs = DL/2)		0.0601	Data Not Lo	gnormal		

#### ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS



Num Obs	Num Miss	Num Valid	Detects	NDs	% NE
Raw Statistics 237	18	219	219	0	0.00%
Number	Minimum	Maximum	Mean	Median	SD
Statistics (Full: no NDs) 219	3.3	257	20.31	17	23.63
K Hat	K Star	Theta Hat	Log Mean	Log Stdv	Log C
Statistics (Full: no NDs) 2.41	2.38	8.427	2.789	0.591	0.212
Normal Distribu	tion Test Res	ults			
No NDs	NDs = DL	NDs = DL/2	Normal ROS	3	
Correlation Coefficient R 0.608	0.608	0.608	0.608		
Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)
Lilliefors (Full: no NDs) 0.29	0.0599	Data Not No	ormal		
Gamma Distribu	ition Test Res	sults			
No NDs	NDs = DL	NDs = DL/2	Gamma RO	S	
Correlation Coefficient R 0.736	0.736	0.736	0.736		
Test value	Crit. (0.05)	5) Conclusion with Alpha(0.05)			5)
Anderson-Darling (Full: no NDs) 8.2	0.764				
Kolmogorov-Smirnov (Full: no NDs) 0.156	0.0621	Data Not Ga	amma Distrib	uted	
Lognormal Distrib	oution Test Re	esults			
No NDs	NDs = DL	NDs = DL/2	Log ROS		
Correlation Coefficient R 0.96	0.96	0.96	0.96		
Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)
Lilliefors (Full: no NDs) 0.103	0.0599	Data Not Lo	gnormal		

## ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS



	Num Obs	Num Miss	Num Valid	Detects	NDs		
Raw Statistics	237	18	219	186	33	15	
	Number	Minimum	Maximum	Mean	Median		
Statistics (Non-Detects Only)	33	3	20	10.18	10	3.0	
Statistics (Detects Only)	186	0.62	157	9.385	3.3	18.	
Statistics (All: NDs treated as DL value)	219	0.62	157	9.505	4.2	17.	
Statistics (All: NDs treated as DL/2 value)	219	0.62	157	8.738	4.2	17.	
Statistics (Normal ROS Estimated Data)	219	-14.19	157	8.647	3.3	17.	
Statistics (Gamma ROS Estimated Data)	219	1E-09	157	9.218	3.5	17.	
Statistics (Lognormal ROS Estimated Data)	219	0.51	157	8.505	3.2	17.	
	K Hat	K Star	Theta Hat	Log Mean	Log Stdv	ı	
Statistics (Detects Only)	0.782	0.774	12	1.478	1.107	0.7	
Statistics (NDs = DL)	0.894	0.885	10.63	1.597	1.067	0.6	
Statistics (NDs = DL/2)	0.87	0.861	10.04	1.493	1.029	0.6	
Statistics (Gamma ROS Estimates)	0.46	0.457	20.02				
Statistics (Lognormal ROS Estimates)				1.41	1.07	0.7	
No	rmal Distribut	tion Test Res	ults				
	No NDs		NDs = DL/2	Normal ROS	3		
Correlation Coefficient R	0.647	0.652	0.629	0.676			
	Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)	
Lilliefors (Detects Only)		0.065	Data Not No		nurr upna(o.o	٠,	
Lilliefors (NDs = DL)		0.0599	Data Not No				
Lilliefors (NDs = DL/2)		0.0599	Data Not No				
Lilliefors (Normal ROS Estimates)		0.0599	Data Not No				
Ga	mma Distribu	tion Test Res	ults				
	No NDs		NDs = DL/2	Gamma RO	S		
Correlation Coefficient R	0.884	0.867	0.859	0.923			
	Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)	
Anderson-Darling (Detects Only)	9.753	0.794					
Kolmogorov-Smirnov (Detects Only)	0.187	0.0699	Data Not Ga	ımma Distribi	uted		
	7.432	0.789					
Anderson-Darling (NDs = DL)			Data Not Gamma Distributed				
Anderson-Darling (NDs = DL) Kolmogorov-Smirnov (NDs = DL)	0.142	0.0635	Data Not Ga	ımma Distribi			
,		0.0635 0.79	Data Not Ga	ımma Distribi			
Kolmogorov-Smirnov (NDs = DL)	10.94			ımma Distribi ımma Distribi	uted		
Kolmogorov-Smirnov (NDs = DL) Anderson-Darling (NDs = DL/2)	10.94 0.221	0.79			uted		
Kolmogorov-Smirnov (NDs = DL) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2)	10.94 0.221 11.6	0.79 0.0636	Data Not Ga				
Kolmogorov-Smirnov (NDs = DL) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (Gamma ROS Estimates) Kolmogorov-Smirnov (Gamma ROS Est.)	10.94 0.221 11.6 0.2	0.79 0.0636 0.831	Data Not Ga	ımma Distribi			
Kolmogorov-Smirnov (NDs = DL) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (Gamma ROS Estimates) Kolmogorov-Smirnov (Gamma ROS Est.)	10.94 0.221 11.6 0.2	0.79 0.0636 0.831 0.0653	Data Not Ga	ımma Distribi			
Kolmogorov-Smirnov (NDs = DL) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (Gamma ROS Estimates) Kolmogorov-Smirnov (Gamma ROS Est.)	10.94 0.221 11.6 0.2 normal Distrib	0.79 0.0636 0.831 0.0653	Data Not Ga Data Not Ga	ımma Distribi			
Kolmogorov-Smirnov (NDs = DL) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (Gamma ROS Estimates) Kolmogorov-Smirnov (Gamma ROS Est.)	10.94 0.221 11.6 0.2 normal Distrib	0.79 0.0636 0.831 0.0653 <b>ution Test Re</b> NDs = DL	Data Not Ga  Data Not Ga	mma Distribu Imma Distribu Log ROS 0.98		5)	
Kolmogorov-Smirnov (NDs = DL) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (Gamma ROS Estimates) Kolmogorov-Smirnov (Gamma ROS Est.)	10.94 0.221 11.6 0.2 normal Distrib No NDs 0.978	0.79 0.0636 0.831 0.0653 <b>ution Test Re</b> NDs = DL 0.985	Data Not Ga  Data Not Ga	umma Distribu Log ROS 0.98  Conclusion w	uted	5)	
Kolmogorov-Smirnov (NDs = DL) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (Gamma ROS Estimates) Kolmogorov-Smirnov (Gamma ROS Est.)  Logr  Correlation Coefficient R	10.94 0.221 11.6 0.2 normal Distrib No NDs 0.978 Test value 0.119	0.79 0.0636 0.831 0.0653 <b>ution Test Re</b> NDs = DL 0.985	Data Not Ga  Data Not Ga  sults  NDs = DL/2 0.982	Log ROS 0.98  Conclusion w	uted	5)	
Kolmogorov-Smirnov (NDs = DL) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (Gamma ROS Estimates) Kolmogorov-Smirnov (Gamma ROS Est.)  Logr  Correlation Coefficient R	10.94 0.221 11.6 0.2 normal Distrib No NDs 0.978 Test value 0.119 0.0924	0.79 0.0636 0.831 0.0653 <b>ution Test Re</b> NDs = DL 0.985 Crit. (0.05) 0.065	Data Not Ga  Data Not Ga  sults  NDs = DL/2 0.982  Data Not Lo	Log ROS 0.98 Conclusion w gnormal gnormal	uted	5)	

## ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS



	Ni Ob	Ni Mina	Nove Malia	Datasta	ND-	0/
Raw Statistics	Num Obs 237	Num Miss 20	Num Valid 217	Detects 61	NDs 156	% 71.
	Number	Minimum	Maximum	Mean	Median	
Statistics (Non-Detects Only)		0.02	0.1	0.0964	0.1	0.01
Statistics (Detects Only)		0.023	0.98	0.119	0.084	0.14
Statistics (All: NDs treated as DL value)		0.02	0.98	0.103	0.1	0.07
Statistics (All: NDs treated as DL/2 value)		0.01	0.98	0.068	0.05	0.08
Statistics (Normal ROS Estimated Data)		-0.228	0.98	0.0402	0.0436	0.11
Statistics (Gamma ROS Estimated Data)		0.023	0.98	0.109	0.0995	0.08
Statistics (Lognormal ROS Estimated Data)		0.0059	0.98	0.0651	0.0456	0.08
		14.0				
0.11.1.15	K Hat	K Star	Theta Hat	Log Mean	Log Stdv	L
Statistics (Detects Only)		1.511	0.0777	-2.492	0.783	-0.3
Statistics (NDs = DL)		3.903	0.026	-2.408	0.503	-0.2
Statistics (NDs = DL/2)		2.413	0.0278	-2.906	0.564	-0.1
Statistics (Gamma ROS Estimates)	3.871	3.821	0.0281			
Statistics (Lognormal ROS Estimates)				-3.09	0.798	-0.2
No	rmal Distribu	tion Test Res	ults			
	No NDs	NDs = DL	NDs = DL/2	Normal ROS	3	
Correlation Coefficient R	0.739	0.59	0.556	0.904		
	Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)
Lilliefors (Detects Only)		0.113	Data Not No		. `	,
Lilliefors (NDs = DL)		0.0601	Data Not No	rmal		
Lilliefors (NDs = DL/2)		0.0601	Data Not No			
Lilliefors (Normal ROS Estimates)		0.0601	Data Not No			
Ga	mma Dietribu	tion Test Res	ulte			
da	No NDs		NDs = DL/2	Gamma RO	S	
Correlation Coefficient R	0.895	0.67	0.706	0.798		
	Test value	Crit. (0.05)		Conclusion w	ith Alpha(0.0	5)
Anderson-Darling (Detects Only)		0.769			. `	,
Kolmogorov-Smirnov (Detects Only)		0.116	Data Not Ga	ımma Distribi	uted	
Anderson-Darling (NDs = DL)		0.757				
Kolmogorov-Smirnov (NDs = DL)		0.062	Data Not Ga	ımma Distribi	uted	
Anderson-Darling (NDs = DL/2)	37.93	0.764				
Kolmogorov-Smirnov (NDs = DL/2)		0.0623	Data Not Ga	ımma Distribi	uted	
Anderson-Darling (Gamma ROS Estimates)		0.757				
Kolmogorov-Smirnov (Gamma ROS Est.)		0.062	Data Not Ga	ımma Distribi	uted	
1	amal Diatrib	ution Toot De	lte			
Logn	No NDs	oution Test Re NDs = DL	NDs = DL/2	Log ROS		
Correlation Coefficient R	0.976	0.82	0.811	0.994		
	Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)
	i cot valac	Ont. (0.00)		r Lognormal	nui / upiia(o.o	0,
Lilliefors (Detects Only)	0.0929	0.113				
Lilliefors (Detects Only)		0.113		•		
Lilliefors (Detects Only) Lilliefors (NDs = DL) Lilliefors (NDs = DL/2)	0.394	0.113 0.0601 0.0601	Data Not Lo	gnormal		

## ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS



	Num Obs	Num Miss	Num Valid	Detects	NDs	9	
Raw Statistics	237	20	217	208	9	4.	
	Number	Minimum	Maximum	Mean	Median		
Statistics (Non-Detects Only)	9	3	4	3.444	3	0.5	
Statistics (Detects Only)	208	2.4	27	9.619	9.4	3.4	
Statistics (All: NDs treated as DL value)	217	2.4	27	9.363	9.3	3.6	
Statistics (All: NDs treated as DL/2 value)	217	1.5	27	9.291	9.3	3.7	
Statistics (Normal ROS Estimated Data)	217	0.503	27	9.32	9.3	3.7	
Statistics (Gamma ROS Estimated Data)	217	1E-09	27	9.318	9.3	3.7	
Statistics (Lognormal ROS Estimated Data)	217	2.4	27	9.389	9.3	3.5	
	K Hat	K Star	Theta Hat	Log Mean	Log Stdv	L	
Statistics (Detects Only)	7.416	7.316	1.297	2.195	0.389	0.1	
Statistics (NDs = DL)	6.251	6.168	1.498	2.155	0.428	0.1	
Statistics (NDs = DL/2)	5.006	4.939	1.856	2.126	0.506	0.2	
Statistics (Gamma ROS Estimates)	2.767	2.732	3.367				
Statistics (Lognormal ROS Estimates)				2.162	0.413	0.1	
No	rmal Distribu	tion Test Res	ults				
	No NDs	NDs = DL	NDs = DL/2	Normal ROS	3		
Correlation Coefficient R	0.972	0.975	0.978	0.979			
	Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)	
Lilliefors (Detects Only)	0.0863	0.0614	Data Not No	rmal			
Lilliefors (NDs = DL)	0.0755	0.0601	Data Not No	rmal			
Lilliefors (NDs = DL/2)	0.0704	0.0601	Data Not Normal				
Lilliefors (Normal ROS Estimates)	0.0724	0.0601	Data Not No	ormal			
Ga	mma Distribu	ition Test Res	sults				
	No NDs	NDs = DL	NDs = DL/2	Gamma RO	S		
Correlation Coefficient R	0.985	0.985	0.978	0.971			
	Test value	Crit. (0.05)	) Conclusion with Alpha(0.05)				
		()	y) Conclusion with Alpha(0.00)				
Anderson-Darling (Detects Only)	1.324	0.754		Conclusion w		5)	
Anderson-Darling (Detects Only) Kolmogorov-Smirnov (Detects Only)				amma Distribi		3)	
• • • • • • • • • • • • • • • • • • • •	0.0651	0.754				٥)	
Kolmogorov-Smirnov (Detects Only)	0.0651 2.044	0.754 0.0628	Data Not Ga		uted	5)	
Kolmogorov-Smirnov (Detects Only) Anderson-Darling (NDs = DL)	0.0651 2.044 0.079	0.754 0.0628 0.755	Data Not Ga	nmma Distribi	uted	5)	
Kolmogorov-Smirnov (Detects Only)  Anderson-Darling (NDs = DL)  Kolmogorov-Smirnov (NDs = DL)	0.0651 2.044 0.079 4.226	0.754 0.0628 0.755 0.0618	Data Not Ga	nmma Distribi	uted	5)	
Kolmogorov-Smirnov (Detects Only)  Anderson-Darling (NDs = DL)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (NDs = DL/2)	0.0651 2.044 0.079 4.226 0.103	0.754 0.0628 0.755 0.0618 0.756	Data Not Ga	amma Distribi amma Distribi	uted	5)	
Kolmogorov-Smirnov (Detects Only)  Anderson-Darling (NDs = DL)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (NDs = DL/2)  Kolmogorov-Smirnov (NDs = DL/2)	0.0651 2.044 0.079 4.226 0.103 11.78	0.754 0.0628 0.755 0.0618 0.756 0.0618	Data Not Ga  Data Not Ga	amma Distribi amma Distribi	uted	5)	
Kolmogorov-Smirnov (Detects Only)  Anderson-Darling (NDs = DL)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (NDs = DL/2)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (Gamma ROS Estimates)  Kolmogorov-Smirnov (Gamma ROS Est.)	0.0651 2.044 0.079 4.226 0.103 11.78 0.183	0.754 0.0628 0.755 0.0618 0.756 0.0618 0.762	Data Not Ga  Data Not Ga  Data Not Ga	amma Distribo amma Distribo amma Distribo	uted	5)	
Kolmogorov-Smirnov (Detects Only)  Anderson-Darling (NDs = DL)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (NDs = DL/2)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (Gamma ROS Estimates)  Kolmogorov-Smirnov (Gamma ROS Est.)	0.0651 2.044 0.079 4.226 0.103 11.78 0.183	0.754 0.0628 0.755 0.0618 0.756 0.0618 0.762 0.0622	Data Not Ga  Data Not Ga  Data Not Ga	amma Distribu amma Distribu amma Distribu	uted	5)	
Kolmogorov-Smirnov (Detects Only)  Anderson-Darling (NDs = DL)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (NDs = DL/2)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (Gamma ROS Estimates)  Kolmogorov-Smirnov (Gamma ROS Est.)	0.0651 2.044 0.079 4.226 0.103 11.78 0.183	0.754 0.0628 0.755 0.0618 0.756 0.0618 0.762 0.0622	Data Not Ga  Data Not Ga  Data Not Ga  Data Not Ga	amma Distribu amma Distribu amma Distribu	uted	5)	
Kolmogorov-Smirnov (Detects Only) Anderson-Darling (NDs = DL) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (Gamma ROS Estimates) Kolmogorov-Smirnov (Gamma ROS Est.)	0.0651 2.044 0.079 4.226 0.103 11.78 0.183	0.754 0.0628 0.755 0.0618 0.756 0.0618 0.762 0.0622 Duttion Test Re	Data Not Ga  Solutis  NDs = DL/2  0.936	amma Distribu amma Distribu amma Distribu amma Distribu Log ROS 0.978	uted		
Kolmogorov-Smirnov (Detects Only) Anderson-Darling (NDs = DL) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (NDs = DL/2) Kolmogorov-Smirnov (NDs = DL/2) Anderson-Darling (Gamma ROS Estimates) Kolmogorov-Smirnov (Gamma ROS Est.)	0.0651 2.044 0.079 4.226 0.103 11.78 0.183 <b>normal Distrib</b> No NDs 0.975	0.754 0.0628 0.755 0.0618 0.756 0.0618 0.762 0.0622 <b>bution Test Re</b> NDs = DL 0.972	Data Not Ga  Solutis  NDs = DL/2  0.936	amma Distribu amma Distribu amma Distribu Log ROS 0.978	uted uted uted		
Kolmogorov-Smirnov (Detects Only)  Anderson-Darling (NDs = DL)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (NDs = DL/2)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (Gamma ROS Estimates)  Kolmogorov-Smirnov (Gamma ROS Est.)  Logr	0.0651 2.044 0.079 4.226 0.103 11.78 0.183 <b>normal Distrib</b> No NDs 0.975 Test value 0.0907	0.754 0.0628 0.755 0.0618 0.756 0.0618 0.762 0.0622 <b>oution Test Re</b> NDs = DL 0.972 Crit. (0.05)	Data Not Ga  NDs = DL/2 0.936	amma Distribu amma Distribu amma Distribu Log ROS 0.978 Conclusion w	uted uted uted		
Kolmogorov-Smirnov (Detects Only)  Anderson-Darling (NDs = DL)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (NDs = DL/2)  Kolmogorov-Smirnov (NDs = DL/2)  Anderson-Darling (Gamma ROS Estimates)  Kolmogorov-Smirnov (Gamma ROS Est.)  Logr  Correlation Coefficient R	0.0651 2.044 0.079 4.226 0.103 11.78 0.183 <b>normal Distrib</b> No NDs 0.975 Test value 0.0907 0.107	0.754 0.0628 0.755 0.0618 0.756 0.0618 0.762 0.0622 <b>oution Test Re</b> NDs = DL 0.972 Crit. (0.05) 0.0614	Data Not Ga  Data Not Ga	amma Distributamma Distributam	uted uted uted		

#### ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS




Num Obs	Num Miss	Num Valid	Detects	NDs	% ND
Raw Statistics 237	20	217	217	0	0.00%
Number	Minimum	Maximum	Mean	Median	SD
Statistics (Full: no NDs) 217	7.6	70	32.46	34	9.579
K Hat	K Star	Theta Hat	Log Mean	Log Stdv	Log C
Statistics (Full: no NDs) 8.983	8.862	3.613	3.423	0.368	0.107
Normal Distribu	tion Test Res	ults			
No NDs	NDs = DL	NDs = DL/2	Normal RO	S	
Correlation Coefficient R 0.976	0.976	0.976	0.976		
Test value	Crit. (0.05)		Conclusion v	vith Alpha(0.0	5)
Lilliefors (Full: no NDs) 0.115	0.0601	Data Not No	ormal		
Gamma Distribe	ition Test Res	sults			
No NDs	NDs = DL	NDs = DL/2	Gamma RO	S	
Correlation Coefficient R 0.952	0.952	0.952	0.952		
Test value	Crit. (0.05)		Conclusion v	vith Alpha(0.0	5)
Anderson-Darling (Full: no NDs) 7.894	0.753				
Kolmogorov-Smirnov (Full: no NDs) 0.169	0.0617	Data Not Ga	amma Distrib	uted	
Lognormal Distril	oution Test Re	esults			
No NDs	NDs = DL	NDs = DL/2	Log ROS		
Correlation Coefficient R 0.924	0.924	0.924	0.924		
Test value	Crit. (0.05)		Conclusion v	vith Alpha(0.0	5)
Lilliefors (Full: no NDs) 0.195	0.0601	Data Not Lo	gnormal		

#### ProUCL 4.00.04 OUTPUT -- GOODNESS-OF-FIT TESTS




_							
Zinc							
		Num Obs	Num Miss	Num Valid	Detects	NDs	% NDs
	Raw Statistics 2	237	20	217	217	0	0.00%
		Number	Minimum			Median	SD
	Statistics (Full: no NDs) 2	217	13.3	607	58.04	51	52.09
		K Hat	K Star	Theta Hat	Log Mean	Log Stdv	Log C\
	Statistics (Full: no NDs) 3	3.501	3.455	16.58	3.912	0.488	0.125
	Norr	nal Distribut	tion Test Res	ults			
		No NDs NDs = DL NDs = DL/2 Normal ROS					
	Correlation Coefficient R (	0.638	0.638	0.638	0.638		
		Test value	Crit. (0.05)	Conclusion with Alpha(0.05)			
	Lilliefors (Full: no NDs) (	).28	0.0601	Data Not Normal			
	Gam	ma Distribu	tion Test Res	ults			
		No NDs	NDs = DL	NDs = DL/2 Gamma ROS			
	Correlation Coefficient R (	).742	0.742	0.742	0.742		
		Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)
	Anderson-Darling (Full: no NDs) 8	3.271	0.759				
	Kolmogorov-Smirnov (Full: no NDs) (	0.166	0.062	Data Not Gamma Distributed			
	Logno	rmal Distrib	ution Test Re	esults			
	•	No NDs	NDs = DL	NDs = DL/2 Log ROS			
	Correlation Coefficient R (	).955	0.955	0.955	0.955		
		Test value	Crit. (0.05)		Conclusion w	vith Alpha(0.0	5)
	Lilliefors (Full: no NDs) (		0.0601	Data Not Lo		,	~,
Motor Cuba	stitution methods such as DL or DL/2 are no			2310 1101 20	g51111d1		
TOTO. GUDS	number metrious such as DE of DE/2 818 IR	A IOCUIIIII	iucu.				


Attachment B-2
Normal Probability Plot for Arsenic




Attachment B-2
Lognormal Probability Plot for Arsenic




Attachment B-2
Normal Probability Plot for Arsenic, Non-detect Concentrations Removed




Attachment B-2
Lognormal Probability Plot for Arsenic, Non-detect Concentrations Removed




Attachment B-2
Normal Probability Plot for Barium




Attachment B-2
Lognormal Probability Plot for Barium




Attachment B-2
Normal Probability Plot for Total Chromium




Attachment B-2
Lognormal Probability Plot for Total Chromium




Attachment B-2
Normal Probability Plot for Cobalt




Attachment B-2
Lognormal Probability Plot for Cobalt




Attachment B-2
Normal Probability Plot for Cobalt, Non-detect Concentrations Removed




Attachment B-2
Lognormal Probability Plot for Cobalt, Non-detect Concentrations Removed




Attachment B-2
Normal Probability Plot for Copper




Attachment B-2
Lognormal Probability Plot for Copper




Attachment B-2
Normal Probability Plot for Lead



Attachment B-2
Lognormal Probability Plot for Lead




Attachment B-2
Normal Probability Plot for Lead, Non-detect Concentrations Removed




Attachment B-2


Lognormal Probability Plot for Lead, Non-detect Concentrations Removed



Attachment B-2
Normal Probability Plot for Mercury




Attachment B-2
Lognormal Probability Plot for Mercury




Attachment B-2

Normal Probability Plot for Mercury, Non-detect Concentrations Removed




Attachment B-2


Lognormal Probability Plot for Mercury, Non-detect Concentrations Removed




Attachment B-2
Normal Probability Plot for Nickel




Attachment B-2
Lognormal Probability Plot for Nickel




Attachment B-2
Normal Probability Plot for Nickel, Non-detect Concentrations Removed




Attachment B-2
Lognormal Probability Plot for Nickel, Non-detect Concentrations Removed




Attachment B-2
Normal Probability Plot for Vanadium




Attachment B-2
Lognormal Probability Plot for Vanadium



Attachment B-2
Normal Probability Plot for Zinc



Attachment B-2
Lognormal Probability Plot for Zinc





Former Pechiney Cast Plate, Inc. Facility Vernon, California

Rosner's	Outlier	lest for	Barium
Rosner's			

Mean92.28Standard Deviation33.06Number of data217Number of suspected outliers1

Potential Critical Obs. Test Critical outlier value (5%) # Mean sd Number value value (1%) 92.28 32.99 190 105 2.962 3.629

For 5% Significance Level, there is no Potential Outlier

For 1% Significance Level, there is no Potential Outlier



Former Pechiney Cast Plate, Inc. Facility Vernon, California

	Rosner's Outlier	Test for	Chromium.	Total
--	------------------	----------	-----------	-------

Mean12.95Standard Deviation4.797Number of data219Number of suspected outliers1

Potential Critical Obs. Test Critical # outlier value (1%) Mean sd Number value value (5%) 12.95 4.786 32.1 27 4.001 3.632 4.003

For 5% Significance Level, there is 1 Potential Outlier Therefore, Observation 32.1 is a Potential Statistical Outlier

For 1% Significance Level, there is no Potential Outlier



Former Pechiney Cast Plate, Inc. Facility Vernon, California


Rosner's Outlier Test for Cobalt

Mean7.951Standard Deviation2.351Number of data203Number of suspected outliers1

Potential Critical Obs. Test Critical outlier value (5%) value (1%) # Mean sd Number value 7.951 2.345 16 129 3.432 3.612 3.984

For 5% Significance Level, there is no Potential Outlier

For 1% Significance Level, there is no Potential Outlier



Former Pechiney Cast Plate, Inc. Facility Vernon, California

Rosner's	Outlier	Test for	Lead
----------	---------	----------	------

Mean9.385Standard Deviation18.7Number of data186Number of suspected outliers1

Critical Potential Obs. Test Critical outlier # Mean sd Number value value (5%) value (1%) 9.385 18.65 157 41 7.915 3.585 3.955

For 5% Significance Level, there is 1 Potential Outlier Therefore, Observation 157 is a Potential Statistical Outlier

For 1% Significance Level, there is 1 Potential Outlier Therefore, Observation 157 is a Potential Statistical Outlier



Former Pechiney Cast Plate, Inc. Facility Vernon, California

Rosner's Outlier Test for Vanadium							
Mean			32.46				
Standard	Deviation		9.579				
Number o	f data		217				
Number o	f suspected	outliers	1				
#	Mean	sd	Potential outlier	Obs. Number	Test value	Critical value (5%)	Critical value (1%)
1	32.46	9.557	70	39	3.928	3.629	4
For 5% Significance Level, there is 1 Potential Outlier Therefore, Observation 70 is a Potential Statistical Outlier							
For 1% Si	ignificance L	evel, there	is no Potenti	al Outlier			